


EXAM II

Physics 101: Lecture 9 Work and Kinetic Energy

- Today's lecture will be on Textbook Sections 6.1 - 6.4



- Previously:



- This is a pain because of vectors...

- There is an easier way to do all this with scalars!

Energy – A Scalar!

- Forms

- Kinetic Energy

Motion (Today)

- Potential Energy

Stored (Next Monday)

- Heat

later

- Mass ($E=mc^2$)

phys 102

- Units: Joules = $\text{kg m}^2 / \text{s}^2 = \text{N m}$

Energy is Conserved

- Energy is “Conserved” meaning it can not be created nor destroyed
 - Can change form
 - Can be transferred
- Total Energy does not change with time.
 - 1. Calculate total energy BEFORE a process
 - 2. Calculate total energy AFTER the process
 - They MUST be equal!!!

This is a BIG deal!

Work: Energy Transfer due to Force

- Force to lift trunk at constant speed to a height d :

➤ Case a $T_a - mg = 0$ $T_a = mg$




» Pull rope down d to move trunk up d

➤ Case b $2T_b - mg = 0$ $T_b = \frac{1}{2} mg$


» Pull rope down d to move trunk up $\frac{1}{2}d$

- $F \times \text{distance}$ must be the same to lift the trunk d

$$\text{Work: } W = F d \cos(\theta)$$



(a)



(b)

Work by Constant Force


A) $W > 0$

B) $W = 0$


C) $W < 0$

- Only component of force parallel to direction of motion does work!

➤ $W = F d \cos \theta$




1)




$W_F < 0: 90 < \theta < 180 : \cos(\theta) < 0$

2)




$W_F = 0: \theta = 90 : \cos(\theta) = 0$

3)



$W_F < 0: 90 < \theta < 180 : \cos(\theta) < 0$

4)



$W_F > 0: 0 < \theta < 90 : \cos(\theta) > 0$

ACTS: Ball Toss

You toss a ball in the air.

What is the work done by gravity as the ball goes up?

- A) Positive B) Negative C) Zero



What is the work done by gravity as the ball goes down?

- A) Positive B) Negative C) Zero

Work by Constant Force

- **Example:** You pull a 30 N chest 5 meters across the floor at a constant speed by applying a force of 50 N at an angle of 30 degrees. How much work is done by the 50 N force?

$$\begin{aligned} W &= F d \cos \theta \\ &= (50 \text{ N}) (5 \text{ m}) \cos (30) \\ &= 217 \text{ J} \end{aligned}$$



Where did the energy go?



- **Example:** You pull a 30 N chest 5 meters across the floor at a constant speed, by applying a force of 50 N at an angle of 30 degrees.
- How much work did gravity do?

$$\begin{aligned} W &= mg d \cos \theta \\ &= 30 * 5 \cos(90) \\ &= 0 \end{aligned}$$

- How much work did friction do?

$$\begin{aligned} \text{X-Direction: } F_{\text{Net}} &= ma \\ T \cos(30) - f &= 0 \\ f &= T \cos(30) \end{aligned}$$

$$\begin{aligned} W &= f d \cos \theta \\ &= 50 \cos(30) * 5 \cos(180) \\ &= -217 \text{ J} \end{aligned}$$




Checkpoint Q1

You are towing a car up a hill with constant velocity. The work done on the car by the normal force is:

- 1. positive
- 2. negative
- 3. zero


← correct



Checkpoint Q2

You are towing a car up a hill with constant velocity. The work done on the car by the gravitational force is:


- 1. positive
- 2. negative ← correct
- 3. zero



Checkpoint Q3

You are towing a car up a hill with constant velocity. The work done on the car by the tension force is:

- 1. positive ← correct
- 2. negative
- 3. zero



Kinetic Energy: Motion

- Apply constant force along x-direction to a point particle m .

$$W = F \Delta x$$

$$= m a \Delta x$$

$$= \frac{1}{2} m (v_f^2 - v_0^2)$$

$$= \frac{1}{2} m v_f^2 - \frac{1}{2} m v_0^2$$

$$\text{recall: } v_f^2 = v_0^2 + 2 a \Delta x$$

$$a \Delta x = \frac{1}{2} (v_f^2 - v_0^2)$$

- Work changes $\frac{1}{2} m v^2$

- Define Kinetic Energy $K = \frac{1}{2} m v^2$

$$W = \Delta K$$


For Point Particles

(i.e. no rotation!)

Checkpoint Q4

You are towing a car up a hill with constant velocity. The total work done on the car by all forces is:

- 1. positive
- 2. negative
- 3. zero ← correct



Example: Block w/ friction

- A block is sliding on a surface with an initial speed of 5 m/s. If the coefficient of kinetic friction μ_k between the block and table is 0.4, how far does the block travel before stopping?

Y direction: $F_{\text{Net}} = ma$

$$N - mg = 0$$

$$N = mg$$

Work

$$W_N = 0$$

$$W_{mg} = 0$$

$$\begin{aligned} W_f &= f d \cos(180) \\ &= -\mu_k m g d \end{aligned}$$


$W = \Delta K$

$$-\mu_k m g d = \frac{1}{2} m (v_f^2 - v_0^2)$$

$$-\mu_k g d = \frac{1}{2} (0 - v_0^2)$$

$$\mu_k g d = \frac{1}{2} v_0^2$$

$$\begin{aligned} d &= \frac{1}{2} v_0^2 / \mu_k g \\ &= 3.1 \text{ meters} \end{aligned}$$



Falling Ball Example

- Ball falls a distance 5 meters, What is final speed?

Only force/work done by gravity

$$W = \Delta K$$

$$W_g = \frac{1}{2} m(v_f^2 - v_i^2)$$

$$F_g d \cos(0) = \frac{1}{2} m v_f^2$$

$$m g d = \frac{1}{2} m v_f^2$$


$$V_f = \text{sqrt}(2 g d) = 10 \text{ m/s}$$



Work by Variable Force



- $W = F_x \Delta x$

- Work is area under F vs x plot



- Spring $F = kx$

- » $\text{Area} = \frac{1}{2} k x^2 = W_{\text{spring}}$



Summary

- Energy is Conserved
- Work = transfer of energy using force
 - Can be positive, negative or zero
 - $W = F d \cos(\theta)$
- Kinetic Energy (Motion)
 - $K = \frac{1}{2} m v^2$
- Work = Change in Kinetic Energy
 - $\Sigma W = \Delta K$