

Last Name: \_\_\_\_\_ First Name \_\_\_\_\_ Network-ID \_\_\_\_\_

Discussion Section: \_\_\_\_\_ Discussion TA Name: \_\_\_\_\_

**Turn off your cell phone and put it out of sight.**

**Keep your calculator on your own desk. Calculators cannot be shared.**

**This is a closed book exam. You have ninety (90) minutes to complete it.**

1. Use a #2 pencil. Do not use a mechanical pencil or pen. Darken each circle completely, but stay within the boundary. If you decide to change an answer, erase vigorously; the scanner sometimes registers incompletely erased marks as intended answers; this can adversely affect your grade. Light marks or marks extending outside the circle may be read improperly by the scanner. Be especially careful that your mark covers the **center** of its circle.

2. You may find the version of **this Exam Booklet at the top of page 2**. Mark the version circle in the TEST FORM box near the middle of your answer sheet. **DO THIS NOW!**

3. Print your **NETWORK ID** in the designated spaces at the *right* side of the answer sheet, starting in the left most column, then **mark the corresponding circle** below each character. If there is a letter "o" in your NetID, be sure to mark the "o" circle and not the circle for the digit zero. If and only if there is a hyphen "-" in your NetID, mark the hyphen circle at the bottom of the column. When you have finished marking the circles corresponding to your NetID, check particularly that you have not marked two circles in any one of the columns.

4. Print **YOUR LAST NAME** in the designated spaces at the *left* side of the answer sheet, then mark the corresponding circle below each letter. Do the same for your **FIRST NAME INITIAL**.

5. Print your UIN# in the STUDENT NUMBER designated spaces and mark the corresponding circles. You need not write in or mark the circles in the SECTION box.

6. Sign your name (**DO NOT PRINT**) on the **STUDENT SIGNATURE line**.

7. On the **SECTION line**, print your **DISCUSSION SECTION**. You need not fill in the COURSE or INSTRUCTOR lines.

*Before starting work, check to make sure that your test booklet is complete. You should have 10 **numbered pages** plus three (3) Formula Sheets following these instructions.*

**Academic Integrity—Giving assistance to or receiving assistance from another student or using unauthorized materials during a University Examination can be grounds for disciplinary action, up to and including dismissal from the University.**

**This Exam Booklet is Version A.** Mark the **A** circle in the **TEST FORM** box near the middle of your answer sheet. **DO THIS NOW!**

*Exam Grading Policy—*

The exam is worth a total of **129** points, composed of three types of questions.

**MC5:** *multiple-choice-five-answer questions, each worth 6 points.*

**Partial credit will be granted as follows.**

- (a) If you mark only one answer and it is the correct answer, you earn **6** points.
- (b) If you mark *two* answers, one of which is the correct answer, you earn **3** points.
- (c) If you mark *three* answers, one of which is the correct answer, you earn **2** points.
- (d) If you mark no answers, or more than *three*, you earn **0** points.

**MC3:** *multiple-choice-three-answer questions, each worth 3 points.*

**No partial credit.**

- (a) If you mark only one answer and it is the correct answer, you earn **3** points.
- (b) If you mark a wrong answer or no answers, you earn **0** points.

**MC2:** *multiple-choice-two-answer questions, each worth 2 points.*

**No partial credit.**


- (a) If you mark only one answer and it is the correct answer, you earn **2** points.
- (b) If you mark the wrong answer or neither answer, you earn **0** points.

Some helpful information:

- A reminder about prefixes: p (pico) =  $10^{-12}$ ; n (nano) =  $10^{-9}$ ;  $\mu$  (micro) =  $10^{-6}$ ; m (milli) =  $10^{-3}$ ; k (kilo) =  $10^3$ ; M or Meg (mega) =  $10^6$ ; G or Gig (giga) =  $10^9$ .


The next two questions pertain to the situation described below.

An uncharged conducting sphere  $A$  hangs from the ceiling by a non-conducting string. As shown in the figure below, a positively charged conducting sphere  $B$  is brought close to the hanging sphere but **does not** touch it.



- 1) What happens to the hanging sphere  $A$ ?
  - a. It moves toward sphere  $B$ .
  - b. It moves away from sphere  $B$ .
  - c. It does not move.
  
- 2) Now the two spheres are **touched** briefly and then separated to the same distance as before. What happens to the hanging sphere  $A$ ?
  - a. It does not move.
  - b. It moves away from sphere  $B$ .
  - c. It moves toward sphere  $B$ .

3) Consider the three molecules to the right. Each atom has a partial charge indicated in the figure. The bond lengths and  $\delta$  are the same in all three molecules.




Which of these molecules has the largest electric dipole moment?

- a. molecule 1
- b. molecule 3
- c. molecule 2

The next three questions pertain to the situation described below.

Consider the following configuration of three point charges  $q_1 = 12 \mu\text{C}$ ,  $q_2 = 5.9 \mu\text{C}$ , and  $q_3 = -2.5 \mu\text{C}$  arranged in a right triangle with sides  $d_{23} = 0.1 \text{ m}$  and  $d_{13} = 3.5 \text{ m}$ . The point charges  $q_2$  and  $q_3$  are both on the x axis of the coordinate system.




- 4) Calculate the x-component  $F_{1tot,x}$  of the force on charge  $q_1$  due to  $q_2$  and  $q_3$ .
- a.  $F_{1tot,x} = -0.0022 \text{ N}$
  - b.  $F_{1tot,x} = 0.003 \text{ N}$
  - c.  $F_{1tot,x} = 0.0015 \text{ N}$
  - d.  $F_{1tot,x} = -0.0018 \text{ N}$
  - e.  $F_{1tot,x} = 6.8 \times 10^{-4} \text{ N}$
- 5) If  $q_1$  were replaced with a charge of the same magnitude but opposite sign,  $-12 \mu\text{C}$ , how would the **magnitude**  $|F_{1tot}|$  of the total force on charge  $q_1$  due to  $q_2$  and  $q_3$  change?
- a. It would increase.
  - b. It would remain the same.
  - c. It would decrease.
- 6) Calculate the magnitude of the electric field,  $E_{tot}$ , at the position of charge  $q_3$  due to charge  $q_1$  and  $q_2$ .
- a.  $E_{tot} = 0 \text{ kV/m}$
  - b.  $E_{tot} = 9100 \text{ kV/m}$
  - c.  $E_{tot} = 7200 \text{ kV/m}$
  - d.  $E_{tot} = 5300 \text{ kV/m}$
  - e.  $E_{tot} = 1.1 \times 10^4 \text{ kV/m}$

7) Consider the following arrangement of three charges on the coordinate axes. All of the charges are positive and have the same magnitude, i.e.,  $Q_1 = Q_2 = Q_3 = +q$ . Note that the grid spacing is the same in x and y.


Where should you place a fourth charge  $Q_4 = +q$  such that the total force on  $Q_3$  is zero, i.e.

$$F_{3tot} = 0?$$



- on the y axis below  $Q_3$ .
- on the y axis above  $Q_3$  but below the origin.
- on the x axis, to the left of  $Q_1$ .
- on the x axis, to the right of  $Q_2$ .
- at the origin.


8) A molecule with an electric dipole moment is placed a distance  $D = 9.9 \text{ nm}$  from an ion of charge  $Q = -e$  and is oriented as shown in the following figure. The molecule has a bond length  $d = 0.1 \text{ nm}$  and partial charges  $\delta = +0.3e$ .



Calculate the magnitude of the total force on the dipole due to the ion.


- $F_{tot} = 1.4 \times 10^{-14} \text{ N}$
- $F_{tot} = 9.3 \times 10^{-15} \text{ N}$
- $F_{tot} = 4 \times 10^{-15} \text{ N}$
- $F_{tot} = 4.5 \times 10^{-14} \text{ N}$
- $F_{tot} = 3.1 \times 10^{-14} \text{ N}$

- 9) A molecule consists of three atoms arranged at an angle  $\theta = 120^\circ$  as shown in the figure below. The atoms have partial charges, with  $\delta = +0.1e$  and the same bond length  $d = 1.9 \times 10^{-10} \text{ m}$ .



Determine the net electric dipole moment  $p$  of the molecule.

- $p = 3 \times 10^{-30} \text{ C}\cdot\text{m}$
  - $p = 2 \times 10^{-30} \text{ C}\cdot\text{m}$
  - $p = 0 \text{ C}\cdot\text{m}$
  - $p = 6.6 \times 10^{-30} \text{ C}\cdot\text{m}$
  - $p = 9.6 \times 10^{-30} \text{ C}\cdot\text{m}$
- 10) Consider a point charge of mass  $m = 19 \text{ g}$  and charge  $q = -6.2 \text{ nC}$  suspended from the ceiling by a massless and non-conducting string of length  $l = 7 \text{ cm}$ . When another point charge of magnitude  $Q = 8 \text{ nC}$  approaches, the hanging charge swings and comes to rest at an angle of  $\theta = 20^\circ$ , as shown in the figure. Assume that the point charge  $q$  comes to rest at the same height as the point charge  $Q$ .




What must be the separation  $d$  between the charges?

- $d = 0.87 \text{ mm}$
- $d = 2.6 \text{ mm}$
- $d = 4.4 \text{ mm}$
- $d = 1.3 \text{ mm}$
- $d = 9.6 \text{ mm}$

The next two questions pertain to the situation described below.

The following figure describes a uniform electric field  $E$  whose magnitude is  $E$ . The dashed lines denote planes perpendicular to the field.




11) What is the electric potential difference  $\Delta V = V_B - V_A$  between points B and A in the figure?

- a.  $\Delta V = \sqrt{2}EL$
- b.  $\Delta V = 0$
- c.  $\Delta V = -\sqrt{2}EL$
- d.  $\Delta V = -EL$
- e.  $\Delta V = EL$

12) A charge  $Q$  is placed initially at B. You drag the charge to point C. What is the work you must do?

- a.  $W = 0$
- b.  $W = -EL$
- c.  $W = EL$


13) What work  $W$  do you have to supply to drag charge  $Q$  from point A along the curve C back to the same point A in the uniform electric field  $E$ ?




- a.  $W < 0$
- b.  $W = 0$
- c.  $W > 0$

14) Which of the following arrangements of equal positive charges has the highest potential energy?

A



B



C




- a. B
- b. C
- c. A



The next two questions pertain to the situation described below.

On the  $y$  axis are two charges  $Q_1 = 1 \mu C$  and  $Q_2 = 3 \mu C$  separated by a distance  $L = 0.4 m$ . Now the charge  $q = 4 \mu C$  is brought to the position A from infinity. (Assume there are no other charges.)




15) What is the required work by you to bring the charge  $q$  to A from infinity?

- a.  $W = 0.44 J$
- b.  $W = 0.17 J$
- c.  $W = 0.15 J$
- d.  $W = 0.22 J$
- e.  $W = 0.65 J$

16) Suppose the work needed by you to bring  $q$  from A to B is  $W$ . If we double both the charges  $Q_1$  and  $Q_2$  the needed work will become


- a.  $W / 2$
- b.  $2 W$
- c.  $4 W$

17) The arrows to the right indicate the electric field emanating from the charge  $q$ . No other charges are present. Point B is in the middle between Point A and Point C. We know Point A has an electric potential of 40 V and Point C has an electric potential of 20 V. What statement about the electric potential  $V_B$  at Point B is correct?




- a.  $V_B < 30 V$
- b.  $V_B = 30 V$
- c.  $V_B > 30 V$

18) Two charges  $Q_1$  and  $Q_2$  are placed on the x-axis with no other charges around. The charge  $Q_1$  is glued to the origin and has a charge of  $q = 3.9 \mu\text{C}$ . The other charge  $Q_2$  is firmly glued to the x-axis at  $x = 5 \text{ m}$ . A small positive test charge  $Q_T$  is brought into the position as shown in the figure and gently released. The test particle does not start to move along the x-axis. What is the value of  $Q_2$ ?



- a.  $Q_2 = 9.8 \mu\text{C}$
- b.  $Q_2 = 24 \mu\text{C}$
- c.  $Q_2 = 1.7 \mu\text{C}$
- d.  $Q_2 = 5.8 \mu\text{C}$
- e.  $Q_2 = 8.8 \mu\text{C}$


19) A charge of  $+|q|$  is placed at the point  $(3, 3)$  and a charge of  $-|q|$  is placed at the point  $(-3, -3)$ . There are no other charges. Which of the following points has a potential of  $0 \text{ V}$ ?



- a.  $(4, -4)$
- b.  $(3, 2)$
- c.  $(4, -5)$
- d.  $(-5, 4)$
- e.  $(-1, -1)$

The next three questions pertain to the situation described below.

Four identical **negative** point charges, each with charge  $-Q$ , are placed at the corners of a square as shown below. The edge length of the square is  $2L$ .



20) What is the magnitude of the electric field  $E$  at the center of the square?

- a.  $E = -2\sqrt{2}k|Q|/L$
- b.  $E = 2\sqrt{2}k|Q|/L^2$
- c.  $E = 0$
- d.  $E = -2\sqrt{2}k|Q|/L^2$
- e.  $E = 2\sqrt{2}k|Q|/L$

21) What is the absolute value of the electric potential at the center of the square?


- a.  $V = 2\sqrt{2}k|Q|/L$
- b.  $V = 8k|Q|/L$
- c.  $V = k|Q|/L$
- d.  $V = 4k|Q|/L^2$
- e.  $V = 0$

22) Now we move the charge  $Q_1$  at the upper right corner of the square to somewhere infinitely far away while keeping the other charges fixed in place. Which of the following statements about the work done **by you** on the charge  $Q_1$  is true?

- a. The work you need to do to move the charge is infinitely large.
- b. The work you need to do to move the charge from the corner of the square to infinity depends on the path we take.
- c. The work you need to do to move the charge is negative.
- d. The work you need to do to move the charge is zero.
- e. The work you need to do to move the charge is positive.

The next three questions pertain to the situation described below.

On a plane there are many point charges. In the following figure eight (8) charges, A, B, C, D, E, F, G, H, and I are depicted with field lines on the plane. The charge A is known to be negative.



23) Choose the correct statement from the following three statements.

- a. B, C, and E are negatively charged.
- b. D, E, and F are positively charged.
- c. F and G are negatively charged.

24) The electric field vanishes at a point very close to


- a. a
- b. c
- c. b

25) Suppose the charge I is allowed to move, but the rest are all fixed. In which direction does charge I start to move?

- a. Roughly in the E direction.
- b. Roughly in the W direction.
- c. Roughly in the S direction.
- d. Roughly in the N direction.
- e. None of N, S, W, E.

The next two questions pertain to the situation described below.

A positive charge  $Q_1 = 6 \mu\text{C}$  is located at position  $(x, y) = (3 \text{ cm}, 3 \text{ cm})$  and a negative charge  $Q_2 = -2 \mu\text{C}$  is located at position  $(x, y) = (-3 \text{ cm}, -3 \text{ cm})$ .



26) What is the potential at the origin of the coordinate system  $(x, y) = (0, 0)$  due to these two charges? No other charges are present.

- a.  $V = 1700 \text{ kV}$
- b.  $V = 850 \text{ kV}$
- c.  $V = 2100 \text{ kV}$
- d.  $V = 340 \text{ kV}$
- e.  $V = 420 \text{ kV}$

27) What is the magnitude of the work done to assemble the charge configuration as shown above?

- a.  $W = 0.52 \text{ J}$
  - b.  $W = 2.6 \text{ J}$
  - c.  $W = 0.65 \text{ J}$
  - d.  $W = 3.2 \text{ J}$
  - e.  $W = 1.3 \text{ J}$
-

# Physic 102 formula sheet (SP2015)

## Kinematics and mechanics

$$x = x_0 + v_0 t + \frac{1}{2} a t^2$$

$$v = v_0 + a t$$

$$v^2 = v_0^2 + 2a\Delta x$$

$$F = ma$$

$$a_c = \frac{v^2}{r}$$

$$E_{\text{tot}} = K + U$$

$$K = \frac{1}{2} m v^2 = \frac{p^2}{2m}$$

$$p = mv$$

$$W_F = F d \cos \theta$$

## Electrostatics

$$F_{12} = k \frac{q_1 q_2}{r^2}$$

$$E = \frac{F}{q_0}$$

$$U_{12} = k \frac{q_1 q_2}{r}$$

$$V \equiv \frac{U}{q_0}$$

$$W_E = -\Delta U = -W_{\text{you}}$$

Point charge

$$E = k \frac{q}{r^2}$$

$$V = k \frac{q}{r}$$

Electric dipole

$$p = qd$$

$$\tau_{\text{dip}} = pE \sin \theta$$

$$U_{\text{dip}} = -pE \cos \theta$$

## Resistance

$$R = \frac{V}{I}$$

$$I = \frac{\Delta q}{\Delta t}$$

$$\text{Physical resistance: } R = \rho \frac{L}{A}$$

$$P = IV = I^2 R = \frac{V^2}{R}$$

$$R_S = R_1 + R_2 + \dots$$

$$\frac{1}{R_P} = \frac{1}{R_1} + \frac{1}{R_2} + \dots$$

## Capacitance

$$C = \frac{Q}{V}$$

$$\text{Parallel plate capacitor: } C = \rho \frac{\kappa \epsilon_0 A}{d}$$

$$E = \frac{Q}{\epsilon_0 A}$$

$$V = Ed$$

$$U_C = \frac{1}{2} QV = \frac{1}{2} CV^2 = \frac{1}{2} \frac{Q^2}{C}$$

$$C_P = C_1 + C_2 + \dots$$

$$\frac{1}{C_S} = \frac{1}{C_1} + \frac{1}{C_2} + \dots$$

## Circuits

$$\sum \Delta V = 0$$

$$\sum I_{\text{in}} = \sum I_{\text{out}}$$

$$q(t) = q_{\infty} (1 - e^{-t/\tau})$$

$$q(t) = q_0 e^{-t/\tau}$$

$$I(t) = I_0 e^{-t/\tau}$$

$$\tau = RC$$

## Magnetism

$$F = qvB \sin \theta$$

$$r = \frac{mv}{qB}$$

$$F_{\text{wire}} = ILB \sin \theta$$

$$\tau_{\text{loop}} = NIAB \sin \varphi$$

Magnetic dipole:

$$\mu = NIA$$

$$\tau_{\text{dip}} = \mu B \sin \varphi$$

$$U_{\text{dip}} = -\mu B \cos \varphi$$

$$B_{\text{wire}} = \frac{\mu_0 I}{2\pi r}$$

$$B_{\text{sol}} = \mu_0 nI$$

## Electromagnetic induction

$$\mathcal{E} = -N \frac{\Delta \Phi}{\Delta t}$$

$$\Phi = BA \cos \varphi$$

$$|\mathcal{E}_{\text{bar}}| = BLv$$

$$\mathcal{E}_{\text{gen}} = \mathcal{E}_{\text{max}} \sin \omega t = \omega NAB \sin \omega t$$

$$\omega = 2\pi f$$

$$V_{\text{rms}} = \frac{V_{\text{max}}}{\sqrt{2}}$$

$$I_{\text{rms}} = \frac{I_{\text{max}}}{\sqrt{2}}$$

$$\frac{V_p}{V_s} = \frac{I_s}{I_p} = \frac{N_p}{N_s}$$

## Electromagnetic waves

$$\lambda = \frac{c}{f}$$

$$E = cB$$

$$u_E = \frac{1}{2} \epsilon_0 E^2$$

$$u_B = \frac{1}{2\mu_0} B^2$$

$$\bar{u} = \frac{1}{2} \epsilon_0 E_{\text{rms}}^2 + \frac{1}{2\mu_0} B_{\text{rms}}^2 = \epsilon_0 E_{\text{rms}}^2 = \frac{B_{\text{rms}}^2}{\mu_0}$$

$$S = I = \bar{u}c = \frac{P}{A}$$

$$f_0 = f_e \sqrt{\frac{1 + v_{\text{rel}}/c}{1 - v_{\text{rel}}/c}} \approx f_e \left(1 + \frac{v_{\text{rel}}}{c}\right)$$

$$I = I_0 \cos^2 \theta$$

## Reflection and refraction

$$\begin{array}{llll} \theta_r = \theta_i & \frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f} & f = \pm \frac{R}{2} & m = \frac{h_i}{h_o} = -\frac{d_i}{d_o} \\ n_1 \sin \theta_1 = n_2 \sin \theta_2 & v = \frac{c}{n} & \sin \theta_c = \frac{n_2}{n_1} & M = \frac{\theta'}{\theta} \approx \frac{d_{\text{near}}}{f} \\ \text{Compound microscope:} & m_{\text{obj}} = \frac{L_{\text{tube}}}{f_{\text{obj}}} & M_{\text{eye}} = \frac{d_{\text{near}}}{f_{\text{eye}}} & M_{\text{tot}} = M_{\text{eye}} m_{\text{obj}} \end{array}$$

## Interference and diffraction

$$\begin{array}{llll} \text{Double-slit interference:} & d \sin \theta = m\lambda & d \sin \theta = \left(m + \frac{1}{2}\right)\lambda & m = 0, \pm 1, \pm 2, \dots \\ \text{Single-slit diffraction:} & a \sin \theta = m\lambda & m = 0, \pm 1, \pm 2, \dots & \\ \text{Circular aperture:} & a \sin \theta \approx 1.22\lambda & & \end{array}$$

## Quantum mechanics

$$\begin{array}{llll} E = hf = \frac{hc}{\lambda} & \lambda = \frac{h}{p} & \Delta p_x \Delta x \geq \frac{\hbar}{2} & \hbar = \frac{h}{2\pi} \\ \text{Bohr atom:} & 2\pi r_n = n\lambda & n = 1, 2, 3, \dots & L_n = m_e v_n r_n = n\hbar \\ r_n = \left(\frac{\hbar^2}{m_e k e^2}\right) \frac{n^2}{Z} \approx (5.29 \times 10^{-11} \text{ m}) \frac{n^2}{Z} & & E_n = -\left(\frac{m_e k^2 e^4}{2\hbar^2}\right) \frac{Z^2}{n^2} \approx -(13.6 \text{ eV}) \frac{Z^2}{n^2} & \\ \frac{1}{\lambda} \approx (1.097 \times 10^7 \text{ m}^{-1}) Z^2 \left(\frac{1}{n_f^2} - \frac{1}{n_i^2}\right) & & & \\ \text{Quantum atom:} & L = \sqrt{\ell(\ell+1)}\hbar & L_Z = m_{\ell}\hbar & S_z = m_s\hbar \\ \text{Atomic magnetism:} & \mu_{e,z} = -\frac{e}{2m_e} L_z & \mu_{s,z} = -\frac{ge}{2m_e} S_z, g \approx 2 & \mu_B \equiv \frac{e\hbar}{2m_e} \approx 5.8 \times 10^{-5} \text{ eV/T} \end{array}$$

## Nuclear physics and radioactive decay

$$\begin{array}{llll} A = Z + N & r \approx (1.2 \times 10^{-15} \text{ m}) A^{1/3} & E_0 = mc^2 & \\ m_{\text{nucleus}} = Zm_{\text{proton}} + Nm_{\text{neutron}} - \frac{|E_{\text{bind}}|}{c^2} & & & \\ \frac{\Delta N}{\Delta t} = \lambda N & N(t) = N_0 e^{-\lambda t} = N_0 2^{-t/T_{1/2}} & T_{1/2} = \frac{\ln 2}{\lambda} \approx \frac{0.693}{\lambda} & \end{array}$$

## Constants and unit conversion

$$\begin{array}{llll} g = 9.8 \text{ m/s}^2 & e = 1.60 \times 10^{-19} \text{ C} & & \\ \epsilon_0 = 8.85 \times 10^{-12} \text{ C}^2/\text{Nm}^2 & k \equiv \frac{1}{4\pi\epsilon_0} = 8.99 \times 10^9 \text{ Nm}^2/\text{C}^2 & \mu_0 = 4\pi \times 10^{-7} \text{ T} \cdot \text{m/A} & \\ c = \frac{1}{\sqrt{\epsilon_0\mu_0}} = 3 \times 10^8 \text{ m/s} & h = 6.626 \times 10^{-34} \text{ J} \cdot \text{s} & hc = 1240 \text{ eV} \cdot \text{nm} & \\ 1 \text{ eV} = 1.60 \times 10^{-19} \text{ J} & m_{\text{electron}} = 9.11 \times 10^{-31} \text{ kg} = 511 \text{ keV}/c^2 & & \\ m_{\text{proton}} = 1.673 \times 10^{-27} \text{ kg} = 938 \text{ MeV}/c^2 & m_{\text{neutron}} = 1.675 \times 10^{-27} \text{ kg} = 939.5 \text{ MeV}/c^2 & & \end{array}$$

## SI Prefixes

| Power             | Prefix | Symbol |
|-------------------|--------|--------|
| 10 <sup>9</sup>   | giga   | G      |
| 10 <sup>6</sup>   | mega   | M      |
| 10 <sup>3</sup>   | kilo   | k      |
| 10 <sup>0</sup>   | —      | —      |
| 10 <sup>-3</sup>  | milli  | m      |
| 10 <sup>-6</sup>  | micro  | $\mu$  |
| 10 <sup>-9</sup>  | nano   | n      |
| 10 <sup>-12</sup> | pico   | p      |