




MATH 241

Midterm 3 Review

Keep in mind that this presentation was created by CARE tutors, and while it is thorough, it is not comprehensive.

QR Code to the Queue




The queue contains the worksheet and the solution to this review session

Fubini's Theorem


- If $f(x,y)$ is continuous on the rectangle

$$R = \{(x,y) \mid a \leq x \leq b, c \leq y \leq d\}$$


$$\iint_R f(x,y) dA = \int_a^b \int_c^d f(x,y) dy dx = \int_c^d \int_a^b f(x,y) dx dy$$



Double Integral Over a General Region



- Integrate dy from $y=x$ to $y=1$
- Then integrate dx



- Integrate dx from $x=0$ to $x=y$
- Then integrate dy

Center of Mass

- The x, y coordinates of the center of mass for an object that has a density function $\rho(x,y)$

$$\bar{x} = \frac{1}{m} \iint x \cdot \rho(x,y) dA \quad \bar{y} = \frac{1}{m} \iint y \cdot \rho(x,y) dA$$

, where mass is calculated as $m = \iint \rho(x,y) dA$

Triple Integral

- Let E be the solid contained under the plane $2x + 3y + z = 6$ in the first octant. Compute the following:

$$\iiint_E 2x \, dV$$

Triple Integral-Cont'd

- Let E be the solid contained under the plane $2x + 3y + z = 6$ in the first octant. Compute the following:







$$\begin{aligned}\iiint_E 2x \, dV &= \int_0^3 \int_0^{2-2x/3} \int_0^{6-2x-3y} 2x \, dz \, dy \, dx = \int_0^3 \int_0^{2-2x/3} 2x(6-2x-3y) \, dy \, dx \\ &= \int_0^3 12x \left(2 - \frac{2x}{3}\right) - 4x^2 \left(2 - \frac{2x}{3}\right) - 3x \left(2 - \frac{2x}{3}\right)^2 \, dx = 9\end{aligned}$$

Example Question #1

- Match the integrals to their corresponding solid regions:

(A) $\int_0^1 \int_y^1 \int_0^{2-x^2-y^2} f(x, y, z) \, dz \, dx \, dy$







(B) $\int_0^1 \int_0^{1-x} \int_0^{1-x^2-y^2} g(x, y, z) \, dz \, dy \, dx$





Example Solution #1

$$(A) \int_0^1 \int_y^1 \int_0^{2-x^2-y^2} f(x, y, z) \, dz \, dx \, dy$$

$$(B) \int_0^1 \int_0^{1-x} \int_0^{1-x^2-y^2} g(x, y, z) \, dz \, dy \, dx$$



base



Polar Coordinates


$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$r^2 = x^2 + y^2$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

$$dA = r dr d\theta$$



<https://magoosh.com/hs/ap-calculus/2017/ap-calculus-bc-review-polar-functions/>

Cylindrical Coordinates

- Cylindrical coordinate is just an extension of polar coordinate to three dimension

$$x = r\cos\theta$$


$$y = r\sin\theta$$

$$z = z$$

$$r^2 = x^2 + y^2$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

$$dV = r dz dr d\theta$$



Sketch of a point in \mathbb{R}^3

Spherical Coordinates


$$x = \rho \sin \varphi \cos \theta$$

$$y = \rho \sin \varphi \sin \theta$$

$$z = \rho \cos \varphi$$

$$\rho^2 = x^2 + y^2 + z^2$$

$$dV = \rho^2 \sin \varphi \, d\rho d\theta d\varphi$$



Sketch of a point in \mathbb{R}^3

Surface Area

- The area of the surface $A(S)$ with equation $z=f(x,y)$ can be calculated as:

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dA$$

Change of Variables Using Jacobian Matrix

- If there is a transformation such that $x=g(u,v)$ and $y=h(u,v)$, then:


$$\iint_R f(x, y) dA = \iint_S f[g(u, v), h(u, v)] \cdot \left| \frac{\partial(x, y)}{\partial(u, v)} \right| d\bar{A}$$

, where the Jacobian Matrix is calculated as

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$


Example Question #2

- Set up the integral to calculate the area of R with the transformation $T(u,v) = (u^2+v, v)$.



Example Solution #2

- Set up the integral to calculate the area of R with the transformation $T(u,v) = (u^2+v, v)$.



$$0 \leq v \leq 1 - \frac{u}{2} \quad 0 \leq u \leq 2$$

Jacobian: $\det \begin{bmatrix} 2u & 1 \\ 0 & 1 \end{bmatrix} = 2u$

Integral: $\int_0^2 \int_0^{1-u/2} 2u \, dv \, du$