



The Grainger College of Engineering

**Center for Academic Resources in Engineering**

---

---

# MATH 241

Midterm 3 Review

---

---

Keep in mind that this presentation was created by CARE tutors, and while it is thorough, it is not comprehensive.

## QR Code to the Queue



The queue contains the worksheet and the solution to this review session


---

# Fubini's Theorem


- If  $f(x,y)$  is continuous on the rectangle

$$R = \{(x,y) \mid a \leq x \leq b, c \leq y \leq d\}$$


$$\iint f(x,y) dA = \int_a^b \int_c^d f(x,y) dy dx = \int_c^d \int_a^b f(x,y) dx dy$$



## Double Integral Over a General Region



- Integrate  $dy$  from  $y=x$  to  $y=1$
- Then integrate  $dx$



- Integrate  $dx$  from  $x=0$  to  $x=y$
- Then integrate  $dy$

# Center of Mass

- The  $x, y$  coordinates of the center of mass for an object that has a density function  $\rho(x, y)$

$$\bar{x} = \frac{1}{m} \iint x \cdot \rho(x, y) dA \quad \bar{y} = \frac{1}{m} \iint y \cdot \rho(x, y) dA$$

, where mass is calculated as  $m = \iint \rho(x, y) dA$

# Triple Integral

- Let  $E$  be the solid contained under the plane  $2x + 3y + z = 6$  in the first octant. Compute the following:

$$\iiint_E 2x \, dV$$

## Triple Integral-Cont'd

- Let  $E$  be the solid contained under the plane  $2x + 3y + z = 6$  in the first octant. Compute the following:







$$\begin{aligned}\iiint_E 2x \, dV &= \int_0^3 \int_0^{2-2x/3} \int_0^{6-2x-3y} 2x \, dz dy dx = \int_0^3 \int_0^{2-2x/3} 2x(6-2x-3y) dy dx \\ &= \int_0^3 12x \left(2 - \frac{2x}{3}\right) - 4x^2 \left(2 - \frac{2x}{3}\right) - 3x \left(2 - \frac{2x}{3}\right)^2 dx = 9\end{aligned}$$

# Example Question #1

- Match the integrals to their corresponding solid regions:

(A)  $\int_0^1 \int_y^1 \int_0^{2-x^2-y^2} f(x, y, z) \, dz \, dx \, dy$

(B)  $\int_0^1 \int_0^{1-x} \int_0^{1-x^2-y^2} g(x, y, z) \, dz \, dy \, dx$









# Example Solution #1

$$(A) \int_0^1 \int_y^1 \int_0^{2-x^2-y^2} f(x, y, z) \, dz \, dx \, dy$$

$$(B) \int_0^1 \int_0^{1-x} \int_0^{1-x^2-y^2} g(x, y, z) \, dz \, dy \, dx$$



base



# Polar Coordinates


$$x = r \cos \theta$$

$$y = r \sin \theta$$

$$r^2 = x^2 + y^2$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

$$dA = r dr d\theta$$



<https://magoosh.com/hs/ap-calculus/2017/ap-calculus-bc-review-polar-functions/>

# Cylindrical Coordinates

- Cylindrical coordinate is just an extension of polar coordinate to three dimension

$$x = r\cos\theta$$


$$y = r\sin\theta$$

$$z = z$$

$$r^2 = x^2 + y^2$$

$$\theta = \arctan\left(\frac{y}{x}\right)$$

$$dV = rdzdrd\theta$$



Sketch of a point in  $\mathbb{R}^3$

# Spherical Coordinates


$$x = \rho \sin\varphi \cos\theta$$

$$y = \rho \sin\varphi \sin\theta$$

$$z = \rho \cos\varphi$$

$$\rho^2 = x^2 + y^2 + z^2$$

$$dV = \rho^2 \sin\varphi \, d\rho d\theta d\varphi$$



Sketch of a point in  $\mathbb{R}^3$

## Surface Area

- The area of the surface  $A(S)$  with equation  $z=f(x,y)$  can be calculated as:

$$A(S) = \iint_D \sqrt{1 + \left(\frac{\partial z}{\partial x}\right)^2 + \left(\frac{\partial z}{\partial y}\right)^2} dA$$

## Change of Variables Using Jacobian Matrix

- If there is a transformation such that  $x=g(u,v)$  and  $y=h(u,v)$ , then:


$$\iint_{\mathbf{R}} f(x, y) dA = \iint_{\mathbf{S}} f[g(u, v), h(u, v)] \cdot \left| \frac{\partial(x, y)}{\partial(u, v)} \right| d\bar{A}$$

, where the Jacobian Matrix is calculated as

$$\frac{\partial(x, y)}{\partial(u, v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \frac{\partial x}{\partial u} \frac{\partial y}{\partial v} - \frac{\partial x}{\partial v} \frac{\partial y}{\partial u}$$


## Example Question #2

- Set up the integral to calculate the area of  $R$  with the transformation  $T(u,v) = (u^2+v, v)$ .



## Example Solution #2

- Set up the integral to calculate the area of  $R$  with the transformation  $T(u,v) = (u^2+v, v)$ .



$$0 \leq v \leq 1 - \frac{u}{2} \quad 0 \leq u \leq 2$$

Jacobian:  $\det \begin{bmatrix} 2u & 1 \\ 0 & 1 \end{bmatrix} = 2u$

Integral:  $\int_0^2 \int_0^{1-u/2} 2u \, dv \, du$