C.A.R.E. PHYS 213 Quiz 1 Review Session


Queue@Illinois
Zaahi (PHYS 213) (zaahimh2)
Logout
CARE / CARE PHYS 213 Exam Review Session
No location specified
Enable notifications
Open Queue
Delete All Questions
Show staff message
This affects anyone using this queue
On-Duty Staff
No on-duty staff
Join
Queue
Welco
sion!
Tutors
uring these times:
Tuesda
Works
Slides
Solutic
Also, h
n the test. If you do not have a Jupyter Notebook environment
installe
y suggestion for this coding example!
Jupyte
Good
This queue is closed. Check back later!

Units for the Exam

- Internal Energy
- Temperature
- Heat Capacity
- Entropy

Internal Energy



- Total energy is **<u>ALWAYS</u>** conserved

- **Positive work on a system increases the system's internal energy**

- **Higher temperature → More Internal Energy**

- First law of thermodynamics:

$$\Delta U = W_{on} + Q$$

Change in internal energy	Work done on the system	Heat added to the system

Temperature & Heat Capacity

- **Heat Capacity** (*C*) - how much energy it takes to **increase** the temperature of a substance by 1 K/°C
  - Units of J/K

- **Larger *C*** $\rightarrow$ **More energy** is required to **increase the temperature** of the object

$$C = dQ/dT$$



Temperature & Heat Capacity

- **Heat Capacity** (*C*) - how much energy it takes to **increase** the temperature of a substance by 1 K/°C
  - Units of J/K

$$C = dQ/dT$$

- **Larger *C*** $\rightarrow$ **More energy** is required to **increase the temperature** of the object

- Water has a larger heat capacity than Aluminum



Types of Heat Capacity

- **Molar Heat Capacity [J/mol K]:** The amount of heat required to raise the temperature of **1 mole** of a substance by 1 K/°C
  - $c_M = C/n$, where n is the number of moles

- **Specific Heat Capacity [J/kg K]:** The amount of heat required to raise the temperature of **1 kg** of a substance by 1 K/°C
  - $c = C/m$, where m is the mass [kg]

- **Heat Capacity at a Constant Volume and Constant Pressure:**
  - $C_v = dU/dT$
  - $C_p = dU/dT + p\, dV/dT$

Equipartition & Heat Capacity

- Only need to memorize DOFs for monatomic gas, diatomic gas, and solids (3, 5, and 6, respectively)

- For substances under the **equipartition assumption:**

$$U = \frac{N_{\text{DOF}}}{2} NkT \implies C = \frac{\mathrm{d}U}{\mathrm{d}T} = \frac{N_{\text{DOF}}}{2} Nk$$

Nk = nR

R = 8.314 J/(mol K)

R = 0.08206 L atm/(mol K)

Equipartition

	Molecule	Degrees of freedom	U	C_V
Monatomic: DOF = 3		$N_{DOF} = 3$ x, y, z momentum	$U = \frac{3}{2}NkT$	$C_V = \frac{3}{2}Nk$
Diatomic: DOF = 5		$N_{DOF} = 5$ x, y, z momentum 2 rotation axes	$U = \frac{(3+2)}{2}NkT$	$C_V = \frac{5}{2}Nk$
Vibrational: DOF = 7		x, y, z momentum 2 rotation axes vibration mode (momentum+potential) $N_{DOF} = 7$	$U = \frac{(3+2+2)}{2}NkT$	$C_V = \frac{7}{2}Nk$
Solid: DOF = 6		x, y, z momentum x, y, z spring modes	$U = \frac{(3+3)}{2}NkT$	$C_V = 3Nk$

Entropy

- Microstate vs Macrostate:
  - Microstate: individual, **specific** arrangement
  - Macrostate: property that arises from the microstates
  - **Many microstates can lead to the same macrostate**
    - Two people have the same weight (macroststate), but the distribution of the weight can be different (microstate)

Entropy (Cont.)

- Entropy (S) is a measure of the degree of 'diversity' associated with a macrostate
  - $S = k \ln(\Omega)$, where Ω is the number of microstates
  - **Second Law of Thermodynamics:** $\Delta S \geq 0$

- Equilibrium
  - Occurs when the macrostate of the system ceases to change
  - The **most probable macrostate** is the one with the **highest entropy (most microstates)**
  - **Equilibrium** is achieved when **S, entropy, is maximized**

Binomial Coefficient

If I have ***N* coins** and I am looking for the macrostate with ***q* heads**, the number of microstates:

$$\binom{N}{q} = \frac{N!}{q!(N-q)!}$$

Example: If I have **20 coins**, how many microstates are associated with the macrostate of getting **7 heads**?

Answer:

Binomial Coefficient

If I have ***N* coins** and I am looking for the macrostate with ***q* heads**, the number of microstates:

$$\binom{N}{q} = \frac{N!}{q!(N-q)!}$$

Example: If I have **20 coins**, how many microstates are associated with the macrostate of getting **7 heads**?

Answer:

$$\binom{20}{7} = \frac{20!}{7!\,(20-7)!} = 77520$$

Entropy (Cont.) - Dice Example

- **Microstate**:
- **Macrostate**:
- What is the most likely macrostate?
  -
  -
- What is the macrostate with the highest entropy?
  -

	⚀	⚁	⚂	⚃	⚄	⚅
⚀	2	3	4	5	6	7
⚁	3	4	5	6	7	8
⚂	4	5	6	7	8	9
⚃	5	6	7	8	9	10
⚄	6	7	8	9	10	11
⚅	7	8	9	10	11	12

- **Entropy of a macrostate is simply a measure of the number of microstates associated with it**
- **More microstates → Higher Entropy, Higher Probability**

Entropy (Cont.) - Dice Example

- **Microstate**: set of individual die values
- **Macrostate**:
- What is the most likely macrostate?
    -
    -
- What is the macrostate with the highest entropy?
    -
- **Entropy of a macrostate is simply a measure of the number of microstates associated with it**
- **More microstates → Higher Entropy, Higher Probability**

	2	3	4	5	6	7
	3	4	5	6	7	8
	4	5	6	7	8	9
	5	6	7	8	9	10
	6	7	8	9	10	11
	7	8	9	10	11	12

Entropy (Cont.) - Dice Example

- **Microstate**: set of individual die values
- **Macrostate**: sum of die values
- What is the most likely macrostate?
  -
  -
- What is the macrostate with the highest entropy?
  -

	⚀	⚁	⚂	⚃	⚄	⚅
⚀	2	3	4	5	6	7
⚁	3	4	5	6	7	8
⚂	4	5	6	7	8	9
⚃	5	6	7	8	9	10
⚄	6	7	8	9	10	11
⚅	7	8	9	10	11	12

- **Entropy of a macrostate is simply a measure of the number of microstates associated with it**
- **More microstates ⟶ Higher Entropy, Higher Probability**

Entropy (Cont.) - Dice Example

- **Microstate**: set of individual die values
- **Macrostate**: sum of die values
- What is the most likely macrostate?
  - 7: has the largest number of microstates associated with it
  - Probability = 6/36 = 0.167

- What is the macrostate with the highest entropy?
  -

- **Entropy of a macrostate is simply a measure of the number of microstates associated with it**
- **More microstates → Higher Entropy, Higher Probability**

	⚀	⚁	⚂	⚃	⚄	⚅
⚀	2	3	4	5	6	7
⚁	3	4	5	6	7	8
⚂	4	5	6	7	8	9
⚃	5	6	7	8	9	10
⚄	6	7	8	9	10	11
⚅	7	8	9	10	11	12

Entropy (Cont.) - Dice Example

- **Microstate**: set of individual die values
- **Macrostate**: sum of die values
- What is the most likely macrostate?
  - 7: has the largest number of microstates associated with it
  - Probability = 6/36 = 0.167

- What is the macrostate with the highest entropy?
  - Again, 7: has the largest number of microstates associated with it

- **Entropy of a macrostate is simply a measure of the number of microstates associated with it**
- **More microstates → Higher Entropy, Higher Probability**

	⚀	⚁	⚂	⚃	⚄	⚅
⚀	2	3	4	5	6	7
⚁	3	4	5	6	7	8
⚂	4	5	6	7	8	9
⚃	5	6	7	8	9	10
⚄	6	7	8	9	10	11
⚅	7	8	9	10	11	12

Important Equations

Start with the definition of temperature

$$\frac{1}{T} \equiv \left(\frac{\partial S}{\partial U}\right)_{V,N}$$

multiply by dU

$$\frac{1}{T}dU = dS$$

integrate

$$\Delta S = \int_{U_i}^{U_f} \frac{1}{T}dU$$

Or the definition of heat capacity

$$C_V = \left(\frac{dQ}{dT}\right)_{V,N}$$

at constant volume
$dQ = dU$

$$C_V = \left(\frac{dU}{dT}\right)_{V,N}$$

multiply by dT

$$C_V dT = dU$$

plug dU into ΔS

$$\Delta S = \int_{T_i}^{T_f} \frac{C_V(T)}{T}dT$$

multiply by dT

$$C_V dT = dQ$$

integrate

$$\Delta U = \int_{T_i}^{T_f} C_V(T)dT$$

Differential Manipulation

- Know these tricks!
- Heat Capacity is the link between *dU* and *dT*
    - **If you know the heat capacity and the temperature change, you can find the change in internal energy and the change in entropy**

$$C = \frac{\partial U}{\partial T} \implies \Delta U = \int_{T_i}^{T_f} C \mathrm{d}T$$

$$\frac{\partial S}{\partial U} = \frac{1}{T} \implies \Delta S = \int \frac{\mathrm{d}U}{T} = \int_{T_i}^{T_f} \frac{C \mathrm{d}T}{T}$$

Entropy of Two Blocks in Thermal Contact - Example

Two blocks A and B, are in thermal contact and insulated from surroundings. Initially, block A is at a higher temperature than block B. **Each block has a temperature-dependent heat capacity given by $C = pT^2$.**

Determine the entropy change for each block



Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

$$C = \frac{dQ}{dT}$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

$$C = \frac{dQ}{dT}$$

$$\Delta S = \int_{T_i}^{T_f} \frac{C\,dT}{T}$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

$$C = \frac{dQ}{dT}$$

$$\boxed{\Delta S = \int_{T_i}^{T_f} \frac{C\,dT}{T}}$$

$$\Delta S = \int_{T_i}^{T_f} \frac{pT^2}{T}\,dT$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

$$C = \frac{dQ}{dT}$$

$$\boxed{\Delta S = \int_{T_i}^{T_f} \frac{C\, dT}{T}}$$

$$\Delta S = \int_{T_i}^{T_f} \frac{pT^2}{T}\, dT$$

$$= \int_{T_i}^{T_f} pT\, dT$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

$$C = \frac{dQ}{dT}$$

$$\boxed{\Delta S = \int_{T_i}^{T_f} \frac{C\,dT}{T}}$$

$$\Delta S = \int_{T_i}^{T_f} \frac{pT^2}{T}\,dT$$

$$= \int_{T_i}^{T_f} pT\,dT$$

$$\boxed{= p\frac{T_f^{\,2}}{2} - p\frac{T_i^{\,2}}{2} = \frac{p}{2}\left[T_f^{\,2} - T_i^{\,2}\right]}$$

Entropy of Two Blocks in Thermal Contact - Example (Cont.)

$$dS = \frac{dQ}{T}$$

$$\Delta S = \int \frac{dQ}{T}$$

$$C = \frac{dQ}{dT}$$

$$\Delta S = \int_{T_i}^{T_f} \frac{C\,dT}{T}$$

$$\Delta S = \int_{T_i}^{T_f} \frac{pT^2}{T}\,dT$$

$$= \int_{T_i}^{T_f} pT\,dT$$

$$= p\frac{T_f^{\,2}}{2} - p\frac{T_i^{\,2}}{2} = \frac{p}{2}\left[T_f^{\,2} - T_i^{\,2}\right]$$

$$\Delta S_A = \frac{p}{2}\left[T_{A,f}^{\,2} - T_{A,i}^{\,2}\right]$$

$$\Delta S_B = \frac{p}{2}\left[T_{B,f}^{\,2} - T_{B,i}^{\,2}\right]$$

Entropy (S) vs. Internal Energy (U)

- Since slope is always positive, temperature is always positive
- **More energy = greater entropy**
- **Diminishing returns:** it gets harder and harder to increase the entropy as internal energy increases
- **Decreasing slope = increasing temperature**
  - **More energy means greater temperature**

$$\frac{1}{T} \equiv \left(\frac{dS}{dU}\right)_{V,N}$$



Good luck!

Feel free to ask any questions you may have.

You got this!