

PHYS 214 Exam 2 Review

CARE / CARE PHYS 214 Exam Review

 No location specified

 Enable notifications



Open Queue

Delete All Questions

Show staff message

This affects anyone using this queue



On-Duty Staff

No on-duty staff

Join

Queue staff message

Welcome to the

Exam Review 1:

- Tuesday, 2

[Worksheet](#)

[Solutions](#)

[Slides](#)

Additionally, he
you can open the

[Jupyter Notebook](#)

Good luck on your



at may help you on your test. If you don't have Jupyter on your computer,

This queue is closed. Check back later!

The queue is empty!

Units Covered on Exam 2

- Unit 4: Photons
- Unit 5: Probability and Complex Numbers
- Unit 6: The Wave Function
- Unit 7: Momentum and Position

Photons

Photons: the quantized bits of light (particles of light)

- Energy of a single photon with frequency f :


$$E = hf = \hbar\omega = \frac{1240 \text{ eV nm}}{\lambda}$$

- Momentum of a single photon with wavelength λ :


$$p = \hbar k = h/\lambda$$

- $h = 6.626 \times 10^{-34} \text{ J} \cdot \text{s}$
- $\hbar$ ('h-bar') = $h/2\pi$

- Photons have a velocity of c :
 - $c = 3 \times 10^8 \text{ m/s}$
 - $c = \lambda f \rightarrow E = (hc)/\lambda$



The Photoelectric Effect




The Photoelectric (Cont.)

This experiment proves the existence of photons and that light can be **BOTH** a particle and a wave

$$KE_{\text{electron}} = eV_{\text{stop}}$$

Stopping Potential:
Voltage applied to stop
electrons from flowing
between the two plates



Work Function (property of the material the light is shining on)

$$KE_{\text{electron}} = hf - \Phi$$

Maximum Kinetic Energy of an ejected electron

Planck's constant times frequency of incoming photons (light)

The diagram shows the equation $KE_{\text{electron}} = hf - \Phi$ enclosed in a hand-drawn cloud-like border. Three arrows point from descriptive text to parts of the equation: one from 'Maximum Kinetic Energy of an ejected electron' to KE_{electron} , one from 'Planck's constant times frequency of incoming photons (light)' to hf , and one from 'Work Function (property of the material the light is shining on)' to Φ .

Effect of Power Source

Increasing the power of a photon source will NOT increase photon energy!

It will only increase photon flux, since frequency/wavelength is what determines photon energy

$$\frac{\# \text{ photons}}{\text{sec}} = \frac{P \text{ Joules}}{\text{sec}} \times \frac{1 \text{ photon}}{X \text{ Joules}}$$

where $X = hf = hc/\lambda$

Force from Photons

$$F = \frac{dp}{dt} = \frac{\text{momentum}}{\text{second}}$$
$$\frac{\text{momentum}}{\text{second}} = \frac{P \text{ joules}}{\text{second}} * \frac{1 \text{ photon}}{hf \text{ joules}} * \frac{\frac{h}{\lambda} \text{ momentum}}{1 \text{ photon}}$$



photons/second

Simplified $\rightarrow F = P/(\lambda f) = P/c$

Example Problem - Photons & Mirrors

A perfectly reflecting mirror is illuminated by a laser beam with a power of **10 mW** and wavelength of **500 nm**.

Determine the force exerted on the mirror by the photons



Example Problem - Photons & Mirrors (Cont.)

$$F = \frac{dp}{dt}$$

Example Problem - Photons & Mirrors (Cont.)

$$F = \frac{dp}{dt}$$

$$F = \frac{2P}{c} = \frac{2(10 * 10^{-3})}{3 * 10^8} = 66.67 \text{ pN}$$

Example Problem - Photons & Mirrors (Cont.)

$$F = \frac{dp}{dt}$$

$$F = \frac{2P}{c} = \frac{2(10 * 10^{-3})}{3 * 10^8} = 66.67 \text{ pN}$$

$$F = 66.67 \text{ pN}$$

Probability and Complex Numbers

Probability

- Probability Density Function:

$$\rho(x)$$

- Probability between two points:

$$P(a \leq x \leq b) = \int_a^b \rho(x) dx$$

- Normalization:

$$\int_{-\infty}^{\infty} \rho(x) dx = 1$$

Complex Numbers

- Two forms:

$$z = a + bi = |z|e^{i\theta}$$

- Conversion:

$$\theta = \arctan(b/a) \quad z = |z|(\cos \theta + i \sin \theta)$$

- Magnitude Squared:

$$|z|^2 = (z^*)(z)$$

- Identities:

$$\cos \theta = \frac{e^{i\theta} + e^{-i\theta}}{2} \quad \sin \theta = \frac{e^{i\theta} - e^{-i\theta}}{2i}$$

The Wave Function

- Wave function is notated as $\Psi(x)$
 - Contains **ALL** information about the properties of a quantum particle
- Properties of $\Psi(x)$:

$$\rho(x) = \psi^*(x)\psi(x) = |\psi(x)|^2$$

$\Psi^*(x)$ is the complex conjugate of $\Psi(x)$

$$P(a \leq x \leq b) = \int_a^b (\Psi^*)(\Psi) dx$$

Probability of detecting a particle
between points a and b

Note: the wave function needs to be normalized

Example Problem - Detection Probability

Given the following normalized wavefunction, what is the integral that gives the chance of measuring the particle in the range $|x| \leq 1$?

$$\Psi(x) = \frac{1}{\sqrt{2}} e^{-|x|/2}$$

Example Problem - Detection Probability (Cont.)

$$P(|x| \leq 1) = \int_{-1}^1 \left(\frac{1}{\sqrt{2}}e^{-|x|/2}\right)^2 dx = \int_0^1 e^{-x} dx \approx 0.63$$

Momentum and Position

- For particles with momentum $\mathbf{p} = \hbar \mathbf{k}$, they are described by this wave function:

$$\psi(x) = Ae^{ikx} \longrightarrow \text{Momentum eigenstate} \\ \text{(1 possible momentum)}$$

- Some particles' wave functions are a superposition of momentum eigenstates:

$$\psi(x) = \boxed{Ae^{ik_1x}} + \boxed{Be^{ik_2x}}$$

1st eigenstate 2nd eigenstate

Momentum and Position (Cont.)

- Probability of each momentum is given by:

$$P(\hbar k_1) = |A|^2 / (|A|^2 + |B|^2)$$


$$P(\hbar k_2) = |B|^2 / (|A|^2 + |B|^2)$$

- Where A and B are the coefficients seen in the wave function: $\psi(x) = \underline{A}e^{ik_1x} + \underline{B}e^{ik_2x}$

- Overall uncertainty is given by **Heisenberg's Uncertainty Principle**:
- The **fewer** momentum eigenstates, the more certain we are for momentum (Δp is small, Δx would be large)
- The **more** momentum eigenstates, the more certain we are for position (Δp is large, Δx is small)


$$\Delta x \Delta p \geq \frac{\hbar}{2}$$

Momentum and Position (cont.)



Momentum ($\rightarrow$ wavelength $\rightarrow$ colour)

$$\Delta x \Delta p \geq \frac{\hbar}{2}$$



Position

Example Problem - Probability of $\hbar k_1$

Determine the probability of the momentum given by $\hbar k_1$:

$$\psi(x) = 2ie^{ik_1x} + 3e^{ik_2x}$$

Example Problem - Probability of $\hbar k_1$ (Cont.)

$$P(\hbar k_1) = \frac{|2i|^2}{|2i|^2 + |3|^2} = \frac{4}{4 + 9} = \frac{4}{13}$$

Good luck on your exam!



Solutions will be posted at the end of the session. Feel free to ask us questions on homework and practice exams as well!

