Exam 5 Review Session
Math 231E



Please join the queue for attendance!

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Outline

1. Please join the queue
2. Mini review of some topics covered
3. Practice! → CARE Worksheet, Practice Exams
   a. Please raise hands for questions rather than put them



Need extra help? → 4th Floor Grainger Library

Subject	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Math 231 (E)	6pm-9pm	12pm-3pm	3pm-8pm		7pm-10pm	1pm-3pm	

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Series / Sequences

- **Sequence:** Just the list of the numbers
  - Limits of sequences
    - Treat it like a function
  - Convergence
    - Treat it like a function
    - Derivative can tell you if it is always increasing or decreasing
- **Series**: The sum of a sequence
  - If a series converges, then the sequence must converge as well.
  - **However:** If sequence converges, then the series may or may not converge.
  - Σa_n converges if the limit of the series converges.

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Integral Test

- Let an = f(n):
  - $\int f(x)dx$ (from k to infinity) converges if the series converges (Σa_k).

    Must be:
    - Continuous
    - Positive
    - Decreasing

1. If $\int_k^\infty f(x)\,dx$ is convergent so is $\sum_{n=k}^{\infty} a_n$.

2. If $\int_k^\infty f(x)\,dx$ is divergent so is $\sum_{n=k}^{\infty} a_n$.

- P-test:
  - The series $\Sigma \frac{1}{n^p}$ converges if p > 1 and diverges if p ≤ 1.

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Comparison/Limit Tests

- **Direct Comparison Test:**
  - Let $0 \leq an \leq bn$.
    - If the series of bn converges, then the series of an does as well.
    - If the series of an diverges, then the series of bn does as well.
- **Limit Comparison Test:**
  - Let $0 \leq an, bn$
    - If the limit of an/bn = C, and C is a nonzero, finite number (ie. not zero or infinity)
    - Then one of two things:
      - Both an and bn converge.
      - Both an and bn diverge.

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Alternating Series Test

- **What is an Alternating Series?**
  - The series is changing signs with each subsequent term
  - $\Sigma a_n(-1)^{n+1}$
- **Alternating Series Test**
  - With series Σa_n, $a_n = (-1)^n b_n$ OR $a_n = (-1)^{n+1} b_n$
    - If $\lim_{n\to\infty} b_n = 0$

      AND
    - b_n is a decreasing sequence
  - **The series Σa_n is convergent**

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Absolute Convergence

- **Absolute Convergence**:
  - If the absolute value of a series, then the series is absolutely convergent.
- **Conditional Convergence**:
  - If a series if convergent, but the absolute value of the series diverges, then the series is conditionally convergent.
- **Negative signs can only help convergence!**

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Root Test

Theorem 2.1 (Root Test). Let a_n be a sequence and $\sum_{n=1}^{\infty} a_n$ be the associated series. Let us define

$$b_n = \sqrt[n]{|a_n|} = |a_n|^{1/n},$$

and assume that $\lim_{n\to\infty} b_n = L$. Then

1. if $L < 1$, then $\displaystyle\sum_{n=1}^{\infty} a_n$ converges absolutely;
2. if $L > 1$, then $\displaystyle\sum_{n=1}^{\infty} a_n$ diverges;
3. if $L = 1$, then no information is obtained.

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Ratio Test

Theorem 3.1. Let a_n be a sequence and $\sum_{n=1}^{\infty} a_n$ be the associated series. Let us assume that

$$\lim_{n\to\infty} \left| \frac{a_{n+1}}{a_n} \right| = L.$$

Then

1. if $L < 1$, then $\sum_{n=1}^{\infty} a_n$ converges absolutely;
2. if $L > 1$, then $\sum_{n=1}^{\infty} a_n$ diverges;
3. if $L = 1$, then no information is obtained.

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Strategies

1. **Check divergence with limit**
2. Look for easy P-Test/Geometric
3. Inspection

TEST	SERIES	CONVERGES IF…	DIVERGES IF…	COMMENTS
*n*th Term Test for Divergence	$\sum_{n=1}^{\infty} a_n$	n/a	$\lim_{n\to\infty} \neq 0$	should be first test used. Inconclusive if limit = 0.
Geometric Series Test	$\sum_{n=1}^{\infty} a_n r^{n-1}$	$\lvert r\rvert < 1$	$\lvert r\rvert \geq 1$	use if there is a "common ratio" $S_n = \frac{a}{1-r}$
P-Series Test	$\sum_{n=1}^{\infty} \frac{1}{n^p}$	$p > 1$	$p \leq 1$	harmonic series when p=1. Useful for comparison tests.
Integral Test	$\sum_{n=1}^{\infty} a_n$ $a_n = f(x)$	$\int_1^{\infty} f(x)\,dx$ converges	$\int_1^{\infty} f(x)\,dx$ diverges	$f(x)$ must be continuous, positive, and decreasing
Direct Comparison Test	$\sum_{n=1}^{\infty} a_n$	$0 \leq a_n \leq b_n$, $\sum_{n=1}^{\infty} b_n$ converges	$0 \leq b_n \leq a_n$, $\sum_{n=1}^{\infty} b_n$ diverges	to show convergence, find a larger series. to show divergence, find a smaller series.
Limit Comparison Test	$\sum_{n=1}^{\infty} a_n$	$\lim_{n\to\infty} \frac{a_n}{b_n} > 0$, $\sum_{n=1}^{\infty} b_n$ converges	$\lim_{n\to\infty} \frac{a_n}{b_n} > 0$, $\sum_{n=1}^{\infty} b_n$ diverges	apply l'hospital's rule if necessary; inconclusive if limit equals 0 or ∞
Alternating Series Test	$\sum_{n=1}^{\infty} (-1)^{n+1} a_n$	$a_{n+1} \leq a_n$, $\lim_{n\to\infty} a_n = 0$	$\lim_{n\to\infty} a_n \neq 0$	must prove that the limit equals 0 and that terms are decreasing
Ratio Test	$\sum_{n=1}^{\infty} a_n$	$\lim_{n\to\infty} \left\lvert\frac{a_{n+1}}{a_n}\right\rvert < 1$	$\lim_{n\to\infty} \left\lvert\frac{a_{n+1}}{a_n}\right\rvert > 1$	test fails if : $\lim_{n\to\infty} \left\lvert\frac{a_{n+1}}{a_n}\right\rvert = 1$
Root Test	$\sum_{n=1}^{\infty} a_n$	$\lim_{n\to\infty} \sqrt[n]{\lvert a_n\rvert} < 1$	$\lim_{n\to\infty} \sqrt[n]{\lvert a_n\rvert} > 1$	test fails if: $\lim_{n\to\infty} \sqrt[n]{\lvert a_n\rvert} = 1$

Power Series

- Can be defined by the form:

$$\sum_{n=0}^{\infty} C_n x^n = C_0 + C_1 x + C_2 x^2 + C_3 x^3 + \cdots$$

  - C_n are the coefficients
  - Series is a function of x
- Can be centered at any number:

$$\sum_{n=0}^{\infty} C_n (x-a)^n = C_0 + C_1 (x-a) + C_2 (x-a)^2 + C_3 (x-a)^3 + \cdots$$

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Power Series

- Domain of Convergence: For what values will the series converge?
  - Use tests to find out what values of x satisfies convergence criteria.

Theorem 3.1. For any power series

$$\sum_{n=0}^{\infty} c_n(x-a)^n,$$

there are exactly three possibilities for the domain of convergence (DOC) and radius of convergence (ROC).

1. Converges only at $x = a$, or
   DOC = $\{a\}$, ROC = 0;
2. Converges for all x, or
   DOC = $(-\infty, \infty)$), ROC = ∞;
3. There is an R such that the power series converges for $|x-a| < R$ and diverges for $|x-a| > R$,
   ROC = R.

Remark 3.2. In the case with a radius of convergence R with $0 < R < \infty$, we have to check the endpoints "by hand".

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN

Good luck on your exam!

You can use the rest of the time to practice

Join the queue!

Subject	Sunday	Monday	Tuesday	Wednesday	Thursday	Friday	Saturday
Math 231 (E)	6pm-9pm	12pm-3pm	3pm-8pm		7pm-10pm	1pm-3pm	

CARE tutoring hours for help (4th Floor, Grainger Library)

UNIVERSITY OF ILLINOIS URBANA-CHAMPAIGN